

2019-05-31

# Southern high-latitude warmth during the JurassicCretaceous: New evidence from clumped isotope thermometry

Vickers, ML

<http://hdl.handle.net/10026.1/14389>

---

10.1130/g46263.1

Geology

Geological Society of America

---

*All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.*

# 1 Southern high latitude warmth during the Jurassic–Cretaceous: 2 New evidence from clumped isotope thermometry

3 **Madeleine L. Vickers<sup>1</sup>, David Bajnai<sup>2</sup>, Gregory D. Price<sup>3</sup>, Jolien Linckens<sup>2</sup>, Jens Fiebig<sup>2</sup>**

4 <sup>1</sup> Department of Geosciences and Natural Resource Management, University of Copenhagen, 1350  
5 Copenhagen, Denmark

6 <sup>2</sup> Institute of Geosciences, Goethe University Frankfurt, 60438 Frankfurt am Main, Germany

7 <sup>3</sup> School of Geography, Earth & Environmental Sciences, University of Plymouth, PL4 8AA  
8 Plymouth, United Kingdom

## 9 **ABSTRACT**

10 In order to understand the climate dynamics of the Mesozoic greenhouse world, it is vital to  
11 determine paleotemperatures from higher latitudes. For the Jurassic and Cretaceous climate, there  
12 are significant discrepancies between different proxies and between proxy data and climate models.  
13 We determined paleotemperatures from Late Jurassic and Early Cretaceous belemnites using the  
14 carbonate clumped isotope paleothermometer and compared these values to temperatures derived  
15 from TEX<sub>86</sub> and other proxies. From our analyses, we infer an average temperature of ca. 25 °C for  
16 the upper part of the water column of the Southern Atlantic Ocean. Our data imply that for mid to  
17 high latitudes, climate models underestimate marine temperatures by >5 °C and, therefore, the  
18 amount of warming that would accompany an increase in atmospheric CO<sub>2</sub> of more than 4x pre-  
19 industrial levels, as is projected for the near future.

## 20 **INTRODUCTION**

21 Modern anthropogenic CO<sub>2</sub> production has resulted in rapid climate change, with near-  
22 surface air temperatures in the high latitude regions rising at ca. twice the global average rate  
23 (Screen and Simmonds, 2010). Predictions of how global and polar temperatures will change over

the next few decades in response to continued CO<sub>2</sub> release may be improved by studying past climate response to elevated CO<sub>2</sub> levels. The Late Jurassic to Early Cretaceous (164 to 100 million years ago) was characterized by extremely high but variable levels of atmospheric CO<sub>2</sub> (from ca. 2x to 8x pre-industrial levels; Wang et al., 2014; Foster et al., 2017), yet reconstructions of marine temperatures, particularly for the high latitudes, are contradictory (e.g., Huber et al., 1995; Price and Gröcke, 2002; Bice et al., 2003; Poulsen, 2004; Jenkyns et al., 2012; Price and Passey, 2013; O'Brien et al., 2017).

The stable oxygen isotope composition of the carbonate remains of marine organisms is the most extensively used temperature proxy, yet high-latitude sea-surface temperatures (SST) derived from independent organic geochemical paleothermometers, i.e., TEX<sub>86</sub>, may be ca. 10 °C warmer than δ<sup>18</sup>O-based temperature reconstructions (e.g., Mutterlose et al., 2010; O'Brien et al., 2012) and up to 6 °C warmer than general circulation model (GCM) predictions (Price and Passey, 2013). These differences have led some authors to suggest that the high TEX<sub>86</sub> SST estimates are too warm (Hollis et al., 2009; Meyer et al., 2018), and there is an ongoing debate as to which calibration is appropriate for applications of the TEX<sub>86</sub> proxy at specific regions and different intervals of the geologic record (Kim et al., 2012; Taylor et al., 2013). Conversely, δ<sup>18</sup>O-based paleotemperature reconstructions rely on several assumptions, among which is the oxygen isotope composition of the seawater (δ<sup>18</sup>O<sub>sw</sub>; Huber et al., 1995; Price and Gröcke, 2002; Bice et al., 2003), and the accuracy of these assumptions still needs to be verified. If the interpretation of warm sub-polar paleo-ocean temperatures can be confirmed, they imply that past and future polar warming may be much greater (i.e., >5 °C) than indicated by climate models. Furthermore, such warm temperatures test the veracity of claims of Early to mid-Cretaceous polar ice, in particular from those studies deriving data from locations distal to the poles (Miller, 2009).

47 Deep Sea Drilling Project Site 511, on the Falkland Plateau (51°00.28'S, 46°58.30'W), is  
48 particularly well suited for studying Jurassic and Cretaceous climate due to its abundant,  
49 exceptionally preserved macrofossils, including belemnites (Jeletzky, 1983; Price and Sellwood,  
50 1997; Price and Gröcke, 2002). The Falkland Plateau was located at approximately 53 °S during the  
51 Late Jurassic–Early Cretaceous (Scotese, 2014; Fig. 1). Mean annual temperatures derived from  
52 GCMs for the Late Jurassic and Early Cretaceous indicate that the temperatures of the Falkland  
53 Plateau region (avg. 10–22 °C) are representative of similar southern hemisphere paleolatitudes  
54 (Lunt et al., 2016). Earlier research at Site 511 used the TEX<sub>86</sub><sup>H</sup> paleotemperature proxy to suggest  
55 that warm sea-surface conditions (26–35 °C) existed during the Late Jurassic–Early Cretaceous  
56 interval (Jenkyns et al., 2012). These paleotemperatures are consistently warmer than  
57 paleotemperature estimates based on  $\delta^{18}\text{O}_{\text{belemnite}}$ , assuming a  $\delta^{18}\text{O}_{\text{sw}}$  of -1‰ SMOW (11–21 °C;  
58 Price and Gröcke, 2002). Another study, undertaken on Barremian to Aptian sediments from two  
59 outcrops in northern Germany, also shows that  $\delta^{18}\text{O}_{\text{belemnite}}$ -derived paleotemperatures (12–16 °C)  
60 are consistently cooler than TEX<sub>86</sub>-based estimates (26–32 °C; Mutterlose et al., 2010). Jenkyns et  
61 al. (2012) argue that the offset is due to TEX<sub>86</sub> recording sea surface temperatures, whereas  
62 belemnites record temperatures from deeper water, possibly from below the thermocline.

63 In this study, we apply the carbonate clumped isotope paleothermometer to exceptionally  
64 well-preserved belemnite rostra from Site 511. This proxy provides seawater temperature estimates  
65 independent of  $\delta^{18}\text{O}_{\text{sw}}$  (Price and Passey, 2013; Wierzbowski et al., 2018). In addition to  
66 constraining high latitude temperatures, we set out to resolve the uncertainties associated with  
67 previous  $\delta^{18}\text{O}$ -based belemnite temperature reconstructions.

## 68 MATERIALS AND METHODS

### 69 Stratigraphy and samples

70 The lithology of the sampled section of Site 511 consists of grey-black, thinly laminated  
71 mudstones and soft, grey claystones, which were deposited in a periodically anoxic, low-energy,  
72 shallow (< 400 m) basin (Basov and Krasheninnikov, 1983; Jeletzky, 1983).

73 A geothermal gradient of 7.4 °C/100 m has been determined (Langseth and Ludwig, 1983)  
74 at Site 511, thus, for the samples analyzed in this study, we can estimate a maximum burial  
75 temperature of ca. 50 °C. At elevated temperatures, diffusion of carbon and oxygen isotopes in the  
76 carbonate mineral lattice may reset the initial bond-ordering (e.g., Henkes et al., 2014). However,  
77 theoretical calculations based on laboratory experiments provide evidence that solid-state diffusion,  
78 even in wet and high-pressure conditions, is insignificant below 100 °C burial temperatures on a  
79 timescale of 100–160 Ma (Passey and Henkes, 2012). Thus, it is unlikely that the belemnite rostra  
80 analyzed in this study were affected by solid-state reordering.

81 Eleven belemnites (*Belemnopsis* sp.) were selected for maximum stratigraphic coverage and  
82 were geochemically screened to include the best-preserved samples, as indicated by available trace  
83 element concentrations (i.e., low Fe and Mn; high Sr and Mg concentrations; Price and Gröcke,  
84 2002; [Supplemental Information](#)) and cathodoluminescence analyses (Figure 5 of Price and  
85 Sellwood, 1997). Subsamples were derived avoiding the margins and apical zone, as these areas are  
86 much more susceptible to diagenetic overgrowth and cementation, respectively than the rest of the  
87 belemnite (e.g., Ullmann et al., 2015). In addition, we made electron backscatter diffraction (EBSD)  
88 analyses and secondary electron microscopy (SEM-BSE) images of selected rostra at the Goethe  
89 University Frankfurt ([Supplemental Information](#)).

## 90 **Clumped Isotope Analyses**

91 Carbonate digestion (90 °C), CO<sub>2</sub> purification (cryotrap and GC) and subsequent  
92 measurement procedures (ThermoFisher MAT 253) are identical to the techniques described in  
93 Wierzbowski et al. (2018). Raw isotope values were calculated using the IUPAC isotopic

parameters, and are projected to the CO<sub>2</sub> reference frame ( $\Delta_{47}^{(RFAC)}$ ; Petersen et al., 2019). To verify the consistency and precision of the clumped isotope measurements, six carbonate standards (ETH1–4, MuStd, Carrara) were analyzed along the samples (Data S1). We used the in-house Wacker et al. (2014) calibration to convert  $\Delta_{47}^{(RFAC)}$  values to temperatures (Supplemental Information; Petersen et al. 2019). Temperature uncertainties are based on external 1SE (including *t*-value) that is always larger than or identical to the best attainable internal precision as represented by the shot noise limit (0.004–0.005‰).

## RESULTS

### Electron Microscopy

All investigated rostra, excluding the areas adjacent to the apical line and the surface, are made up of optical calcite and the c-axis of the calcite grains point radially outwards (Figs. S1-S4). The distribution of the crystallographic a-axes also follows a pattern. This is analogous to pristinely preserved rostra (Stevens et al., 2017). Our EBSD and SEM-BSE analyses suggest that recrystallization, which would change the original orientation of the biogenic calcite grains, did not occur in the sampled areas.

### Clumped Isotope Analyses

The  $\Delta_{47}^{(RFAC)}$  values range between 0.690(±0.011)‰ and 0.707(±0.015)‰. The 1SE uncertainty for the clumped isotope measurements, calculated from 4–6 replicate analyses are between 0.004‰ and 0.015‰ (mean 0.010‰). The  $\Delta_{47}^{(RFAC)}$  values yield seawater temperatures ranging between 21 °C and 28 °C (mean 25 °C) and show no significant stratigraphic trend (Fig. 2). The average uncertainty for the reconstructed temperatures is ±4 °C. Steeper-sloped calibrations yield indistinguishable temperatures within ±1SE (Data S1).

## DISCUSSION

117       The  $\Delta_{47}$ -derived temperature range (21–28 °C, mean 25 °C) for the entire section is higher  
118       than those temperatures reconstructed via stable oxygen isotope paleothermometry (11–19 °C,  
119       mean 16 °C, assuming  $\delta^{18}\text{O}_{\text{sw}} = -1\text{‰}$  SMOW; Price and Gröcke, 2002), and cooler, and rarely  
120       within error, of SST estimates derived from TEX<sub>86</sub> (25–31 °C; Fig. 2; Jenkyns et al., 2012). In this  
121       study, as in Jenkyns et al. (2012), we calculate TEX<sub>86</sub> temperatures using the TEX<sub>86</sub><sup>H</sup> calibration  
122       (Kim et al., 2010). Given the shallow-water and high latitude setting of Site 511 TEX<sub>86</sub><sup>H</sup> may yield  
123       maximum SST estimates (Schouten et al., 2013; Taylor et al., 2013). In contrast to TEX<sub>86</sub><sup>H</sup>, the  
124       linear calibration used of O'Brien et al. (2017) yield ca. 2–3 °C warmer temperatures, whereas the  
125       calibrations that assume a non-surface export depth of GDGTs (Kim et al., 2012; Schouten et al.,  
126       2013) yield ca. 5–6 °C cooler estimates (Fig. 2). Although the TEX<sub>86</sub><sup>H</sup> proxy is likely the most  
127       appropriate for a high latitude setting such as Site 511, there is ongoing discussion and revision of  
128       the various calibrations, and ongoing debate as to which calibration should be applied (e.g., Ho et  
129       al., 2014; Inglis et al., 2015). The difference between the TEX<sub>86</sub><sup>H</sup> and the  $\Delta_{47}$ -derived temperatures  
130       for Site 511 may be partially resolved by considering a seasonal bias in either proxy. It has been  
131       postulated that belemnites, as nektonic cephalopods, reflect mean annual temperatures (MAT; Price  
132       and Sellwood, 1997; Mutterlose et al., 2010), while TEX<sub>86</sub> may indicate summer temperatures,  
133       rather than MAT (Leider et al., 2010; Hollis et al., 2012). Nevertheless, our  $\Delta_{47}$  temperatures  
134       suggest that belemnites were calcifying their rostra in the upper part of the water column (<200 m  
135       depth), and are broadly consistent with TEX<sub>86</sub>-derived SSTs, given the uncertainties listed above.  
136       Such an interpretation is in alignment with an assumed predator lifestyle in the photic zone for  
137       belemnites (Klug et al., 2016).

138       All three records from Site 511 show less than 7 °C variability across the entire Late  
139       Jurassic and Early Cretaceous interval, although the low sampling resolution means it is not  
140       possible to derive more detailed information on Jurassic and Cretaceous climate evolution. These

141 data confirm warm Late Jurassic–Early Cretaceous high latitude ocean temperatures, possibly  
142 precluding the likelihood of substantial land ice, and are consistent with estimated MATs from  
143 fossil plant assemblages from the Antarctic Peninsula (Francis and Poole, 2002). The most likely  
144 mechanism to account for such warmth observed at Site 511 is high atmospheric greenhouse gas  
145 concentrations and high polar heat transport. The shallow meridional temperature gradients of the  
146 past greenhouse climates pose a significant challenge to numerical climate models (Huber and  
147 Caballero, 2011), in that increased greenhouse gases may yield warm Polar Regions, but also  
148 overheat the Tropics. MATs for the Cretaceous derived from coupled ocean-atmosphere climate  
149 models provide estimates for 53 °S ranging from 12 °C to 21 °C (Zhou et al., 2008; Donnadieu et  
150 al., 2016). The higher of these estimates are generated with 2240 ppm  $p\text{CO}_2$  (8 x pre-industrial  
151 levels; Donnadieu et al., 2016). These atmospheric  $\text{CO}_2$  concentrations typically exceed estimates  
152 of Cretaceous  $p\text{CO}_2$  derived from fossil leaf stomatal index measurements, isotope-based or  
153 geochemical model estimates (Wang et al., 2014; Foster et al., 2017).

154 Furthermore, it is crucial to consider the magnitude of a non- $\text{CO}_2$  component of local  
155 climate change, before proxies from a single site are interpreted in a global context (Lunt et al.,  
156 2016). GCM output indicates warm conditions during the Cretaceous at Site 511 when compared to  
157 the Eocene (Lunt et al., 2016), with almost invariable modeled global mean temperatures over the  
158 same period, when  $p\text{CO}_2$  is kept constant. This suggests that contributions from other processes  
159 (e.g., paleogeography) may account for some of the observed warmth. Despite these findings and  
160 those of others (Donnadieu et al., 2016), the role of paleogeography in regulating climate remains  
161 less than clear.

162 Such warm temperatures at Site 511 challenge our understanding of how the ocean-  
163 atmosphere system operated in the past (Poulsen, 2004) and may also have important implications  
164 for the prediction of future climates as they imply we may be underestimating future climate change



165 in such regions (Spicer et al., 2008). Proposed mechanisms to increase the transfer of heat toward  
166 the poles (Schmidt and Mysak, 1996), including sensible and latent heat transfer via the atmosphere  
167 and heat transfer via the oceans (Hotinski and Toggweiler, 2003), are hence implied. As Site 511  
168 was situated in a seaway open to the southwest (Fig. 1), increased heat transfer via warm ocean  
169 currents can only be derived from the Pacific. Thus, other processes, including heat transfer via the  
170 atmosphere, might also be important for this region.

171 These new warm  $\Delta_{47}$ -derived temperature reconstructions also have implications for basin-  
172 scale hydrologies. In conjunction with the  $\delta^{18}\text{O}_{\text{belemnite}}$  data (Price and Sellwood, 1997; Price and  
173 Gröcke, 2002), we can estimate  $\delta^{18}\text{O}_{\text{sw}}$ , assuming the temperature dependence of oxygen isotope  
174 fractionation between belemnite calcite and seawater corresponds to Kim and O'Neil (1997). The  
175  $\delta^{18}\text{O}$ -temperature equation of Kim and O'Neil (1997) indicates that  $\delta^{18}\text{O}_{\text{sw}}$  may have averaged  
176  $+1.0\text{‰}$  SMOW ( $1SE = 0.7\text{‰}$ ; Fig. 2, Data S1), heavier than the global average for an ice-free world  
177 ( $-1\text{‰}$  SMOW; Shackleton and Kennett, 1975). This could suggest that the semi-enclosed basin in  
178 which Site 511 was located was dominated by evaporation; alternatively, it is quite possible that the  
179 Kim and O'Neil (1997) calcite equation is not applicable to belemnite calcite.

## 180 CONCLUSIONS

181 This proxy-to-proxy intercomparison reduces the uncertainty on temperature estimates for  
182 the Mesozoic high southern latitudes. Our  $\Delta_{47}$ -derived temperatures, although slightly cooler, are  
183 consistent with the  $\text{TEX}_{86}^{\text{H}}$  reconstructions for sea-surface temperatures. The new  $\Delta_{47}$  data, in  
184 conjunction with  $\delta^{18}\text{O}_{\text{belemnite}}$  data imply local  $\delta^{18}\text{O}_{\text{sw}}$  values of ca.  $1.0(\pm 0.7)\text{‰}$  SMOW, indicating a  
185 strong role of evaporation on the Falkland Plateau, which was a semi-enclosed basin during the Late  
186 Jurassic and Early Cretaceous. The warm reconstructed paleotemperatures, if extrapolated  
187 poleward, reinforce evidence of temperate polar conditions and lack of polar ice. If these warm  
188 ocean temperatures, occurring when  $p\text{CO}_2$  in Earth's atmosphere were also high, prove accurate,

189 they may indicate that greenhouse gases could have heated the oceans during the Jurassic and  
190 Cretaceous more than currently accepted. This suggests that future warming from elevated  
191 atmospheric CO<sub>2</sub> concentrations may be much greater than that predicted by models.

## 192 **ACKNOWLEDGMENTS**

193 We thank C. John (Imperial College, London), S. Hofmann, C. Schreiber (Goethe  
194 University Frankfurt), N. Löffler, K. Methner, E. Krsnik (Senckenberg BIK-F), D. Gröcke (Durham  
195 University), S. Robinson (University of Oxford) and our anonymous reviewers. Funding was  
196 provided by a PhD scholarship from the University of Plymouth, UK, and a European Consortium  
197 for Ocean Research Drilling (ECORD) grant to M.V.; a UK Natural Environment Research Council  
198 (NERC) grant (NE/J020842/1) to G.D.P.; and the European Union's Horizon 2020 research and  
199 innovation program under the Marie Skłodowska-Curie grant agreement No. 643084 (BASE-LiNE  
200 Earth) to B.D and J.F.

## 201 **REFERENCES CITED**

- 202 Basov, I. A., and Krasheninnikov, V., 1983, Benthic foraminifers in Mesozoic and Cenozoic  
203 sediments of the southwestern Atlantic as an indicator of paleoenvironment, Deep Sea  
204 Drilling Project Leg 71: Initial Reports of the Deep Sea Drilling Project, v. 71, p. 739-787.
- 205 Bice, K. L., Huber, B. T., and Norris, R. D., 2003, Extreme polar warmth during the Cretaceous  
206 greenhouse? Paradox of the late Turonian  $\delta^{18}\text{O}$  record at Deep Sea Drilling Project Site 511:  
207 Paleooceanography, v. 18, no. 2.
- 208 Donnadieu, Y., Puc  at, E., Moiroud, M., Guillocheau, F., and Deconinck, J.-F., 2016, A better-  
209 ventilated ocean triggered by Late Cretaceous changes in continental configuration: Nature  
210 communications, v. 7, p. 10316.
- 211 Foster, G. L., Royer, D. L., and Lunt, D. J., 2017, Future climate forcing potentially without  
212 precedent in the last 420 million years: Nature Communications, v. 8, p. 14845.

213 Francis, J. E., and Poole, I., 2002, Cretaceous and early Tertiary climates of Antarctica: evidence  
 214 from fossil wood: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 182, no. 1, p. 47-  
 215 64.

216 Henkes, G. A., Passey, B. H., Grossman, E. L., Shenton, B. J., Pérez-Huerta, A., and Yancey, T. E.,  
 217 2014, Temperature limits for preservation of primary calcite clumped isotope  
 218 paleotemperatures: *Geochimica et cosmochimica acta*, v. 139, p. 362-382.

219 Ho, S. L., Mollenhauer, G., Fietz, S., Martínez-García, A., Lamy, F., Rueda, G., Schipper, K.,  
 220 Méheust, M., Rosell-Melé, A., and Stein, R., 2014, Appraisal of TEX<sub>86</sub> and TEX<sub>86</sub><sup>L</sup>  
 221 thermometries in subpolar and polar regions: *Geochimica et Cosmochimica Acta*, v. 131, p.  
 222 213-226.

223 Hollis, C. J., Handley, L., Crouch, E. M., Morgans, H. E., Baker, J. A., Creech, J., Collins, K. S.,  
 224 Gibbs, S. J., Huber, M., and Schouten, S., 2009, Tropical sea temperatures in the high-  
 225 latitude South Pacific during the Eocene: *Geology*, v. 37, no. 2, p. 99-102.

226 Hollis, C. J., Taylor, K. W., Handley, L., Pancost, R. D., Huber, M., Creech, J. B., Hines, B. R.,  
 227 Crouch, E. M., Morgans, H. E., and Crampton, J. S., 2012, Early Paleogene temperature  
 228 history of the Southwest Pacific Ocean: Reconciling proxies and models: *Earth and*  
 229 *Planetary Science Letters*, v. 349, p. 53-66.

230 Hotinski, R., and Toggweiler, J., 2003, Impact of a Tethyan circumglobal passage on ocean heat  
 231 transport and “equable” climates: *Paleoceanography*, v. 18, no. 1.

232 Huber, B. T., Hodell, D. A., and Hamilton, C. A., 1995, Mid to Late Cretaceous climate of the  
 233 southern high latitudes: stable isotopic evidence for minimal equator-to-pole thermal  
 234 gradients: *Bulletin of the Geological Society of America*, v. 107, p. 1164–1191.

235 Huber, M., and Caballero, R., 2003, Eocene El Niño: Evidence for robust tropical dynamics in the  
 236 “hothouse”: *Science*, v. 299, no. 5608, p. 877-881.

237 Inglis, G. N., Farnsworth, A., Lunt, D., Foster, G. L., Hollis, C. J., Pagani, M., Jardine, P. E.,  
 238 Pearson, P. N., Markwick, P., and Galsworthy, A. M., 2015, Descent toward the Icehouse:  
 239 Eocene sea surface cooling inferred from GDGT distributions: *Paleoceanography and*  
 240 *Paleoclimatology*, v. 30, no. 7, p. 1000-1020.

241 Jeletzky, J., 1983, Macroinvertebrate paleontology, biochronology, and paleoenvironments of  
 242 Lower Cretaceous and Upper Jurassic rocks, Deep Sea Drilling Hole 511, eastern Falkland  
 243 Plateau: *Initial Reports of the Deep Sea Drilling Project*, v. 71, p. 951-975.

244 Jenkyns, H., Schouten-Huibers, L., Schouten, S., and Sinninghe Damsté, J., 2012, Warm Middle  
 245 Jurassic-Early Cretaceous high-latitude sea-surface temperatures from the Southern Ocean:  
 246 *Climate of the Past*, v. 8, no. 1.

247 Kim, J.-H., Romero, O. E., Lohmann, G., Donner, B., Laepple, T., Haam, E., and Damste, J. S. S.,  
 248 2012, Pronounced subsurface cooling of North Atlantic waters off Northwest Africa during  
 249 Dansgaard–Oeschger interstadials: *Earth and Planetary Science Letters*, v. 339, p. 95-102.

250 Kim, J.-H., Van der Meer, J., Schouten, S., Helmke, P., Willmott, V., Sangiorgi, F., Koç, N.,  
 251 Hopmans, E. C., and Damsté, J. S. S., 2010, New indices and calibrations derived from the  
 252 distribution of crenarchaeal isoprenoid tetraether lipids: Implications for past sea surface  
 253 temperature reconstructions: *Geochimica et Cosmochimica Acta*, v. 74, no. 16, p. 4639-  
 254 4654.

255 Kim, S.-T., and O'Neil, J. R., 1997, Equilibrium and nonequilibrium oxygen isotope effects in  
 256 synthetic carbonates: *Geochimica et Cosmochimica Acta*, v. 61, no. 16, p. 3461-3475.

257 Klug, C., Schweigert, G., Fuchs, D., Kruta, I., and Tischlinger, H., 2016, Adaptations to squid-style  
 258 high-speed swimming in Jurassic belemnitids: *Biology letters*, v. 12, no. 1, p. 20150877.

259 Langseth, M., and Ludwig, W., 1983, A heat-flow measurement on the Falkland Plateau: *Initial*  
 260 *Reports of the Deep Sea Drilling Project*, v. 71, no. SEP, p. 299-303.

261 Leider, A., Hinrichs, K.-U., Mollenhauer, G., and Versteegh, G. J., 2010, Core-top calibration of the  
 262 lipid-based  $U^{K'}_{37}$  and TEX<sub>86</sub> temperature proxies on the southern Italian shelf (SW Adriatic  
 263 Sea, Gulf of Taranto): *Earth and Planetary Science Letters*, v. 300, no. 1-2, p. 112-124.

264 Lunt, D. J., Foster, G. L., O'Brien, C. L., Pancost, R. D., and Robinson, S. A., 2016,  
 265 Palaeogeographic controls on climate and proxy interpretation: *Climate of the Past*, v. 12,  
 266 no. 5, p. 1181.

267 Meyer, K. W., Petersen, S. V., Lohmann, K. C., and Winkelstern, I. Z., 2018, Climate of the Late  
 268 Cretaceous North American Gulf and Atlantic Coasts: *Cretaceous Research*, v. 89, p. 160-  
 269 173.

270 Miller, K. G., 2009, Broken greenhouse windows *Nature Geoscience*, v. 2, p. 465-466.

271 Mutterlose, J., Malkoc, M., Schouten, S., Sinninghe Damsté, J. S., and Forster, A., 2010, TEX<sub>86</sub> and  
 272 stable  $\delta^{18}O$  paleothermometry of early Cretaceous sediments: Implications for belemnite  
 273 ecology and paleotemperature proxy application: *Earth and Planetary Science Letters*, v.  
 274 298, no. 3, p. 286-298.

275 O'Brien, C. L., Robinson, S. A., Pancost, R. D., Damsté, J. S. S., Schouten, S., Lunt, D. J., Alsenz,  
 276 H., Bornemann, A., Bottini, C., and Brassell, S. C., 2017, Cretaceous sea-surface  
 277 temperature evolution: Constraints from TEX<sub>86</sub> and planktonic foraminiferal oxygen  
 278 isotopes: *Earth-Science Reviews*, v. 172, p. 224-247.

279 Passey, B. H., and Henkes, G. A., 2012, Carbonate clumped isotope bond reordering and  
 280 geospeedometry: *Earth and Planetary Science Letters*, v. 351, p. 223-236.

281 Petersen, S. V., Defliese, W. F., Saenger, C., Daëron, M., Huntington, K. W., John, C. M., Kelson,  
 282 J. R., Coleman, A. S., Kluge, T., Olack, G. A., Schauer, A. J., Bajnai, D., Bonifacie, M.,  
 283 Breitenbach, S. F., Fiebig, J., Fernandez, A. B., Henkes, G. A., Hodell, D., Katz, A., Kele, S.,  
 284 Lohmann, K. C., Passey, B. H., Peral, M. Y., Petrizzo, D. A., Rosenheim, B. E., Tripathi, A.,

285 Venturelli, R., Young, E. D., and Winkelstern, I. Z. 2019, Effects of improved <sup>17</sup>O correction  
286 on inter-laboratory agreement in clumped isotope calibrations, estimates of mineral-specific  
287 offsets, and acid fractionation factor temperature dependence: *Geochemistry, Geophysics,*  
288 *Geosystems*.

289 Poulsen, C. J., 2004, Palaeoclimate: a balmy Arctic: *Nature*, v. 432, no. 7019, p. 814-815.

290 Price, G.D., and Sellwood, B., 1997, “Warm” palaeotemperatures from high Late Jurassic  
291 palaeolatitudes (Falkland Plateau): Ecological, environmental or diagenetic controls?:  
292 *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 129, no. 3, p. 315-327.

293 Price, G. D., and Gröcke, D. R., 2002, Strontium-isotope stratigraphy and oxygen-and carbon-  
294 isotope variation during the Middle Jurassic–Early Cretaceous of the Falkland Plateau,  
295 South Atlantic: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 183, no. 3, p. 209-  
296 222.

297 Price, G. D., and Passey, B. H., 2013, Dynamic polar climates in a greenhouse world: Evidence  
298 from clumped isotope thermometry of Early Cretaceous belemnites: *Geology*, v. 41, p. 923-  
299 926.

300 Schmidt, G. A., and Mysak, L. A., 1996, Can increased poleward oceanic heat flux explain the  
301 warm Cretaceous climate?: *Paleoceanography*, v. 11, no. 5, p. 579-593.

302 Schouten, S., Hopmans, E. C., and Damsté, J. S. S., 2013, The organic geochemistry of glycerol  
303 dialkyl glycerol tetraether lipids: a review: *Organic geochemistry*, v. 54, p. 19-61.

304 Scotese, C. R., 2014, Atlas of Early Cretaceous Paleogeographic Maps, PALEOMAP Atlas for  
305 ArcGIS, volume 2, The Cretaceous, Maps 23-31, Mollweide Projection, PALEOMAP  
306 Project, Evanston, IL.

307 Screen, J. A., and Simmonds, I., 2010, The central role of diminishing sea ice in recent Arctic  
308 temperature amplification: *Nature*, v. 464, no. 7293, p. 1334-1337.

309 Shackleton, N., and Kennett, J., 1975, Paleotemperature history of the Cenozoic and the initiation of  
310 Antarctic glaciation: oxygen and carbon isotope analyses in DSDP Sites 277, 279, and 281,  
311 Washington DC, US Government Printing Office, Initial reports of the deep sea drilling  
312 project, 743-755 p.:

313 Spicer, R. A., Ahlberg, A., Herman, A. B., Hofmann, C.-C., Raikevich, M., Valdes, P. J., and  
314 Markwick, P. J., 2008, The Late Cretaceous continental interior of Siberia: A challenge for  
315 climate models: *Earth and Planetary Science Letters*, v. 267, no. 1, p. 228-235.

316 Stevens, K., Griesshaber, E., Schmahl, W., Casella, L. A., Iba, Y., and Mutterlose, J., 2017,  
317 Belemnite biomineralization, development, and geochemistry: The complex rostrum of  
318 *Neohibolites minimus*: *Palaeogeography, palaeoclimatology, palaeoecology*, v. 468, p. 388-  
319 402.

320 Taylor, K. W., Huber, M., Hollis, C. J., Hernandez-Sanchez, M. T., and Pancost, R. D., 2013, Re-  
321 evaluating modern and Palaeogene GDGT distributions: Implications for SST  
322 reconstructions: *Global and Planetary Change*, v. 108, p. 158-174.

323 Ullmann, C., Frei, R., Korte, C., and Hesselbo, S. P., 2015, Chemical and isotopic architecture of  
324 the belemnite rostrum: *Geochimica et Cosmochimica Acta*, v. 159, p. 231-243.

325 Wacker, U., Fiebig, J., Tödter, J., Schöne, B. R., Bahr, A., Friedrich, O., Tütken, T., Gischler, E.,  
326 and Joachimski, M. M., 2014, Empirical calibration of the clumped isotope  
327 paleothermometer using calcites of various origins: *Geochimica et Cosmochimica Acta*, v.  
328 141, p. 127-144.

329 Wang, Y., Huang, C., Sun, B., Quan, C., Wu, J., and Lin, Z., 2014, Paleo-CO<sub>2</sub> variation trends and  
330 the Cretaceous greenhouse climate: *Earth-Science Reviews*, v. 129, p. 136-147.

331 Wierzbowski, H., Bajnai, D., Wacker, U., Rogov, M. A., Fiebig, J., and Tesakova, E. M., 2018,  
332 Clumped isotope record of salinity variations in the Subboreal Province at the Middle–Late  
333 Jurassic transition: Global and Planetary Change.

334 Zhou, J., Poulsen, C., Pollard, D., and White, T., 2008, Simulation of modern and middle  
335 Cretaceous marine  $\delta^{18}\text{O}$  with an ocean-atmosphere general circulation model:  
336 Paleooceanography, v. 23, no. 3, p. PA3223.

337



338

339 **FIGURE CAPTIONS**

340 **Figure 1.** Paleogeographic setting of the Deep Sea Drilling Project Site 511. Early Cretaceous  
341 paleogeographic reconstruction after Scotese (2014).



**Figure 2.** Jurassic and Early Cretaceous temperatures and seawater  $\delta^{18}\text{O}$  from DSDP Site 511. **(A)** Clumped isotope seawater temperature reconstructions for Site 511 (this study) are compared to those based on  $\delta^{18}\text{O}_{\text{belemnite}}$  (Price and Gröcke, 2002; Price and Sellwood, 1997; plotted using Kim and O’Neil, 1997, with an assumed  $\delta^{18}\text{O}_{\text{sw}}$  of -1‰ SMOW) and  $\text{TEX}_{86}$  (Jenkyns et al., 2012). Infilled green circles represent  $\delta^{18}\text{O}_{\text{belemnite}}$  temperatures from Price and Gröcke (2002), hollow green circles are the belemnites that were also used for clumped isotopes analysis in this study. For  $\text{TEX}_{86}$  temperatures, dotted lines used  $\text{TEX}_{86}^{\text{H},0-200}$  (Kim et al. 2012., eq. 2), dashed used  $\text{TEX}_{86}\text{-linear}$  (O’Brien et al., 2017, eq. 4), and solid line and points used the  $\text{TEX}_{86}^{\text{H}}$  calibration (Kim et al. 2010, eq.10). **(B)** Reconstructed  $\delta^{18}\text{O}_{\text{sw}}$  values (this study) using the equation of Kim and O’Neil (1997). Error bars represent for  $\delta^{18}\text{O}_{\text{belemnite}}$  and  $\Delta_{47}$  the 1SE of multiple replicate analyses; for  $\text{TEX}_{86}^{\text{H}}$  the calibration error; and for  $\delta^{18}\text{O}_{\text{sw}}$  the 1SE. corresponding to the  $\Delta_{47}$  measurements. Age model construction is described in the [Supplemental Information](#), whereas data for this figure can be found in [Data S1](#).

